

# Msx3-iCre

This line is useful for driving Cre-mediated recombination of reporter lines, or for deleting floxed genes in the dorsal spinal cord neuroepithelium.

To generate this line, we used a 135Kb mouse genomic PAC (PAC 402-G09) isolated from library RCPI21 (Oseogawa et al., 2000). The library contains genomic DNA from a female mouse, strain 129S6/SvEvTac1.

The DNA region used included all Msx3 coding exons as well as  $\sim$ 64Kb upstream and  $\sim$ 68Kb downstream (of the Msx3 gene). We inserted the codon-improved version of Cre recombinase (iCre) (Shimshek et al., 2002) followed by an SV40 polyadenylation signal into the first coding exon of Msx3, fusing iCre to the endogenous translation initiation codon.

The modified PAC was linearized by AscI digestion prior to generating transgenic mice by pronuclear injection. In the developing mouse spinal cord, transgene expression can be detected in the dorsal half of the spinal cord neuroepithelium (Fig.1).

Generation and characterization of the mouse has been described (Fogarty, 2005; Tripathi et al., 2011). The mouse is also described here: <a href="https://www.informatics.jax.org/allele/MGI:5430776">https://www.informatics.jax.org/allele/MGI:5430776</a>

# Msx3 iCre

Fig. 1:  $\frac{\ln situ}{\ln situ}$  hybridization showing expression of Msx3 and iCre in the embryonic mouse spinal cord at E11.5.

### **Category**

Biological Materials/Genetically Modified Organisms

## Placing an order on XIP

To license this product, please select the **appropriate licence option** on the right-hand side of this page. Terms can be previewed from the "Preview terms" link.

MTAs require agreement between all the parties involved in supplying and receiving a product. This cannot happen instantaneously but is a controlled process, managed through XIP and should not take longer no than 10 business days in ordinary circumstances.

To place an order, please locate the <u>Sign-in</u> or <u>Register</u> options on the top right side of this page. You can either sign in to your existing account or register for a new now.

For additional guidance on how to create an account and place an order, refer to the <u>FAQs</u>. Please note that your account should be created using your academic/ institutional email address.

### References

- D R Shimshek, J Kim, M R Hübner, D J Spergel, F Buchholz, E Casanova, A F Stewart, P H Seeburg, R Sprengel(08 January 2002), https://onlinelibrary.wiley.com/journal/1526968x, https://onlinelibrary.wiley.com/doi/abs/10.1002/gene.10023, 32, 19-26
- Kazutoyo Osoegawa, Minako Tateno, Peng Yeong Woon, Eirik Frengen, Aaron G. Mammoser, Joseph J. Catanese, Yoshihide Hayashizaki, and Pieter J. de Jong(January 2000), https://www.ncbi.nlm.nih.gov/pmc/articles/PMC310499/, https://genome.cshlp.org/, 10, 116-128
- 3. Fogarty MH(January 2006), https://discovery.ucl.ac.uk/id/eprint/1445454/, 1-235
- 4. Tripathi RB; Clarke LE; Burzomato V; Kessaris N; Anderson PN; Attwell D; Richardson WD(4 May 2011) , https://pubmed.ncbi.nlm.nih.gov/21543611/, https://www.jneurosci.org/, 31, 6809-6819